

Simplifications of the Tensors in Quantum Scattering, Part I

Gordon Chalmers

e-mail: gordon@quartz.shango.com

Abstract

In this comment it is pointed out that the perturbative dynamics of general massive field theories can be mapped to delimited sums of determined integral functions. The limits on the sums are the remaining obstacle to finding the general g -loop, or derivative expanded, form of the scattering.

In previous works the classical scattering of scalar and gauge field theories has been formulated with the use of partitions of numbers [1],[2]. The use of the classical scattering has been used in the formulation of the perturbative quantum dynamics [3],[4]. There are simplifications in the quantum scattering with the use of the determined integral functions, presented in [3]. The content of this letter is to show that the perturbative dynamics of a general massive theory can be reduced to the, as yet delimited, sums of numbers parameterizing the products of these integral functions.

In massive ϕ^3 theory, with coupling g , the scattering takes on the form [3],

$$g^L m^q \prod s_{ij}^{n_{ij}} \sum \prod_i B(n_i, m_i, w_i) , \quad (1)$$

with the parameters n_i , m_i and w_i . There are conditions on the sum which depend on the loop order L and the kinematic configuration of n_{ij} . The B functions are integral moments defined and calculated in [2].

The simplest example is symmetry breaking in QED,

$$\mathcal{L} = -\frac{1}{4}F^2 + \frac{1}{2}\nabla\phi^\dagger\nabla\phi + \bar{\psi}^a\gamma^\mu\nabla_\mu\psi_a , \quad (2)$$

with $\nabla_\mu = \partial_\mu + gA_\mu$; the masses of the fermion and scalar are left non-zero. The vacuum value of the scalar is chosen as,

$$\nabla\phi^\dagger\nabla\phi = \partial\phi^\dagger\partial\phi + 2gA(v+\phi)^*\partial\phi + c.c. + g|v+\phi|^2A^2 , \quad (3)$$

which gives a mass to the gauge field.

The non-abelian gauge theory is defined by,

$$\mathcal{L} = -\frac{1}{4}F^b F_b + \frac{1}{2}\nabla\phi^a\nabla\phi_a \quad (4)$$

with the covariant derivatives,

$$\nabla = \partial + gA \quad \nabla_\mu\phi^a = \partial_\mu\phi_a + g[\phi, A_\mu]_a . \quad (5)$$

The symmetry breaking could be achieved as in the standard $SU(N) \rightarrow U(1)$, with additional interactions to give masses to the abelian fields.

One example of symmetry breaking is as follows,

$$\phi_a \rightarrow \phi_a + v_a \quad (6)$$

with,

$$\partial\phi^a\partial\phi_a + g^2 f^{cab} f_c^{de} (v_a + \phi_a)(v_d + \phi_d) A_b A_e \quad (7)$$

$$+ g 2 \text{Tr} T_c [T_a, T_b] (v_b + \phi_b) A_c \partial^\mu \phi^a . \quad (8)$$

The mass term is

$$(v_a + \phi_a)(v^a + \phi^a) A_b A^b - [(v_a + \phi_a) A^a]^2 \quad (9)$$

The vacuum expectation values are chosen uniformly $v_a = v\eta_a$ ($\eta^2 = 1$), so that

$$v^2 A^2 - v^2 \eta \cdot A^2 = v^2 [A^2 - (\sum \eta_a A_a)^2] \quad (10)$$

$$= v^2 \sum (1 - \eta_a^2) A_a^2 - 2v^2 \sum \eta_a \eta_b A_a A_b , \quad (11)$$

which can be used to give mass to all of the the gauge fields. The gauge fixing term could be written in axial gauge as $v^2(\eta \cdot A)^2$, and makes the uniform breaking manifest.

In three dimensions, a gauge invariant mass could be generated with the topological term

$$m^2 F \tilde{F} , \quad (12)$$

without any use of the scalar fields.

The gauge scattering, with n external gauge field states, then takes on the form,

$$\prod_{i=1}^m \varepsilon_i \cdot \rho_j \ g^q \ m_A^{p_1} m_\phi^{p_2} \prod s_{ij}^{n_{ij}} \sum \prod_i B(n_i, m_i, w_i) , \quad (13)$$

with the possible tensors

$$\rho_j = \epsilon_j \quad \text{or} \quad k_j . \quad (14)$$

The complete quantum gauge scattering then is produced by the appropriate summations on the indices in the sum in (13). This expansion depends on the helicity configuration in (14), that is choosing the contractions of momenta or helicities as ρ_i , with m ranging from $n/2$ to n (all polarizations or all momenta as ρ_i).

In the example of a three dimensional gauge field with the mass term, the form is simpler,

$$\prod_{i=1}^m \varepsilon_i \cdot \rho_j \ g^L m^p \prod s_{ij}^{n_{ij}} \sum \prod_i B(n_i, m_i, w_i) , \quad (15)$$

with the indices on the sum reflecting the dynamics of the three dimensional massive gauge theory. The summations on the B functions depend on the helicity configuration and the powers of the coupling.

In any theory with the mass generation for the gauge fields, and with non-zero values for the remaining particle content, there is an expansion of the form (1), (13), or (15), with the appropriate mass values used in their forms. The integral functions $B(n_i, m_i, w_i)$ are determined. The completion of the quantum scattering is determined by the appropriate summations of the indices. The required non-analyticity due to unitarity, such as $\ln(s - m^2)$ is found by resumming the polynomial terms.

These summations on the indices are potentially determined through a simple context, such as the use of some analog of the Hopf algebra associated with the tree diagrams. The determination of the sum limits in the scattering determines the perturbative quantum scattering, with possible redundancy in the sums, and in a general massive theory with various particle content.

References

- [1] Gordon Chalmers, *Tree Amplitudes in Scalar Field Theories*, physics/0504173.
- [2] Gordon Chalmers, *Tree Amplitudes in Gauge and Gravity Theories*, physics/0504219.
- [3] Gordon Chalmers, *Quantum Solution to Scalar Field Theory Models*, physics/0505018.
- [4] Gordon Chalmers, *Quantum Gauge Theory Amplitude Solutions*, physics/0505077.

Tensor Simplifications, Part II: Amplitude Parameterizations

Gordon Chalmers

e-mail: gordon@quartz.shango.com

Abstract

The amplitudes in gauge theory, gravity, and gauge theories with matter are parameterized. The coefficients of the parameterized amplitudes are required in order to specify the complete coupling form, and in determining the couplings within the derivative expansion. However, these parameterizations can be used in a similar manner as a Fourier transform in a fit to experimental data over large energy ranges.

The computation of gauge theory amplitudes is typically complicated due to both the loop integrals and the tensor computations. Many tools have been developed to circumvent these difficulties. The computation of gauge theory amplitudes has been extensively studied for many years. The parameterization of the amplitudes is useful in their computation within the derivative expansion [1]-[11]. The general form of the gauge theory amplitudes has uses in particle phenomenology.

The parameterization of the amplitudes has never been elucidated in a clear context in the literature. The general form of the the pure gauge theory amplitude is

$$\prod \varepsilon_{\sigma(i)} \cdot \varepsilon_{\tilde{\sigma}(j)} \prod \varepsilon_{\rho(i)} \cdot k_{\tilde{\rho}(j)} \quad (16)$$

$$\frac{\prod t_{a(i)}^{\tilde{a}(i)}}{\prod t_{b(i)}^{\tilde{b}(i)}} \prod \ln^{d_1(i)} \dots \ln^{d_m(i)} t_{c(i)}^{\tilde{c}(i)} . \quad (17)$$

The sets $\sigma, \tilde{\sigma}, \rho, \tilde{\rho}, a, \tilde{a}, b, \tilde{b}, c, \tilde{c}, d_l$ are required in order to specify the general form of the amplitude. In the case of massive theories there are factors of the mass parameters. The inclusion of the fermions requires the factors

$$\prod \psi_a^\alpha \psi_{b\alpha} \prod \bar{\psi}_a^{\dot{\alpha}} \bar{\psi}_{b\dot{\alpha}} \prod \psi_i^\alpha k_{\alpha\kappa(i)} \bar{\psi}^{\dot{\alpha},i} k_{\dot{\alpha}\tilde{\kappa}(i)} \prod \psi_i^\alpha \bar{\psi}_j^{\dot{\alpha}} \varepsilon_{\alpha\dot{\alpha}}(\nu(i)) , \quad (18)$$

which extends from the external line factors.

There is a further group theory tensor,

$$\prod_j \text{Tr} \prod_i T_{\gamma_j(i)} , \quad (19)$$

with as many traces there are loops in the amplitude calculation. The pure gravity amplitude, also in perturbation theory, requires the same form, except there are no traces; there is a doubling of the ε 's into holomorphic and non-holomorphic parts, ε_i and $\bar{\varepsilon}_i$.

There is a numerical factor multiplying the tensor form, and is proportional to the gauge coupling, g^{n-2+2L} . This numerical coefficient is required to specify the complete amplitude, and at each loop order.

The numerical factor

$$f(g, M) = \sum b_L(M) g^{n-2+2L} , \quad (20)$$

with M representing the set of values σ , ρ , a , $\tilde{a}$, b , $\tilde{b}$, c , and $\tilde{c}$, d_l . These coefficients can be iteratively determined in the gauge theory derivative expansion. There is a restriction on the number of logarithms due to the number of loops. Possible functional forms of the scattering are,

$$\rho_M \sum \dots \sum F_{p,q} \dots F_{p,q} , \quad (21)$$

for the gauge scattering, and termwise for the graviton scattering (which has a dimensionful coupling constant),

$$\gamma_M \Gamma \dots \Gamma / \Gamma \dots \Gamma . \quad (22)$$

These forms are line with lower dimensional field theories.

However, despite the determination of the amplitudes in a straightforward calculational scheme, their form may be used to model partonic cross-sections; the linear combination of the terms in (17) can be used to define cross-sections in the usual manner. In other words, a large set of these terms in (17) can be used to model a greater range in energies in particle amplitudes used in, for example, QCD or electroweak cross sections. Their form, as a collective set, can model very large ranges in energies, even without knowing the explicit coupling dependence in (20).

An interesting further question is the inversion of a general linear combination of the terms in (17) into an effective Lagrangian and its properties.

References

- [1] G. Chalmers, *Quantum Scalar Field Theory Solution*, physics/0505018.
- [2] G. Chalmers, *Derivation of Quantum Field Dynamics*, physics/0503062.
- [3] G. Chalmers, *Quantum Gauge Theory Solutions*, physics/0505077.
- [4] G. Chalmers, *Masses and Interactions of Nucleons in the Derivative Expansion*, physics/0503110.
- [5] G. Chalmers, *$N=4$ Supersymmetric Gauge Theory in the Derivative Expansion*, hep-th/0209088.
- [6] G. Chalmers, *Gauge Theories in the Derivative Expansion*, hep-th/0209086.
- [7] G. Chalmers, *Scalar Field Theory in the Derivative Expansion*, hep-th/0209075.
- [8] G. Chalmers, *Quantum Gravity with the Standard Model*, hep-th/0209072.
- [9] G. Chalmers, *On the Finiteness of $N=8$ Supergravity*, hep-th/0008162.
- [10] G. Chalmers and J. Erdmenger, *Dual expansions of $N=4$ super Yang-Mills theory via IIB superstring theory*, Nucl. Phys. B **585**:517 (2000), hep-th/0005192.
- [11] G. Chalmers, *S and U -duality Constraints on IIB S -matrices*, Nucl. Phys. B **580**:193 (2000), hep-th/0001190.